

# A Note on Groups with Just-Infinite Automorphism Groups

**Francesco de Giovanni**

*Dipartimento di Matematica e Applicazioni, Università di Napoli Federico II, via Cintia,  
I - 80126 Napoli (Italy) degiovan@unina.it*

**Diana Imperatore**

*Dipartimento di Matematica e Applicazioni, Università di Napoli Federico II, via Cintia,  
I - 80126 Napoli (Italy) diana.imperatore@unina.it*

Received: 6.9.2012; accepted: 5.10.2012.

**Abstract.** An infinite group is said to be *just-infinite* if all its proper homomorphic images are finite. We investigate the structure of groups whose full automorphism group is just-infinite.

**Keywords:** just-infinite group; automorphism group

**MSC 2000 classification:** 20E36

## 1 Introduction

Let  $D_\infty$  be the infinite dihedral group. Although  $D_\infty$  admits non-inner automorphisms, it is well known that  $D_\infty$  is isomorphic to its full automorphism group  $\text{Aut}(D_\infty)$ . Moreover, it has been proved in [3] that there are no other groups with infinite dihedral automorphism group. In other words, the equation

$$\text{Aut}(X) \simeq D_\infty$$

admits, up to isomorphisms, only the trivial solution  $X = D_\infty$ . It is usually hard to understand which groups  $Q$  can occur as full automorphism groups of some other group, i.e. when the equation

$$\text{Aut}(X) \simeq Q$$

admits at least one solution. For instance, it was proved by D.J.S. Robinson [8] that no infinite Černikov group can be realized as full automorphism group of a group, while a classical result of R. Baer [1] showed that for any finite group  $Q$  the above equation has no solution within the universe of infinite periodic groups.

The aim of this paper is to obtain informations about groups whose automorphism groups are just-infinite. Here a group  $G$  is said to be *just-infinite* if it is infinite but all its proper homomorphic images are finite. It follows from Zorn's Lemma that any finitely generated infinite group has a just-infinite homomorphic image, and so just-infinite groups play a relevant role in many problems of the theory of infinite groups (see for instance [4]). The first examples of just-infinite groups are of course the infinite cyclic group and the infinite dihedral group; the first of these cannot occur as the full automorphism group of any group, and we discussed above the dihedral case. We will prove that if  $G$  is any group admitting an ascending normal series whose factors are either central or finite, then the automorphism group  $\text{Aut}(G)$  cannot be just-infinite. It follows in particular that in any group with just-infinite automorphism group, the centre and the hypercentre coincide. Some examples of groups with just-infinite automorphism groups will also be constructed.

Most of our notation is standard and can for instance be found in [7].

## 2 Results and examples

The structure of just-infinite groups has been described by J.S. Wilson [11]; of course, all infinite simple groups are just-infinite, while any soluble-by-finite just-infinite group is a finite extension of a free abelian group of finite rank. In the same paper, the class  $\mathfrak{D}_2$ , consisting of all infinite groups in which every non-trivial subnormal subgroup has finite index, is considered; obviously, all  $\mathfrak{D}_2$ -groups are just-infinite, but it is clear that any  $\mathfrak{D}_2$ -group containing an abelian non-trivial subnormal subgroup is either cyclic or dihedral. It follows easily from this remark that the result proved in [3] can be extended to the next statement (recall here that a group is called *generalized soluble* if it has an ascending subnormal series whose factors are either abelian or finite).

**Theorem 1.** *Let  $G$  be a generalized soluble group whose automorphism group  $\text{Aut}(G)$  is a  $\mathfrak{D}_2$ -group. Then  $G \simeq D_\infty$ .*

The following example shows that a similar result cannot be proved for groups with just-infinite automorphism groups, even restricting the attention to the case of polycyclic groups.

Let  $A = \langle a \rangle \times \langle b \rangle$  be a free abelian group of rank 2, and let  $x$  and  $y$  be the automorphisms of  $A$  defined by the positions

$$a^x = b, \quad b^x = a, \quad a^y = b, \quad b^y = a^{-1}b.$$

Then  $\langle x, y \rangle$  is a dihedral subgroup of order 12 of  $GL(2, \mathbb{Z})$ , and the semidirect product

$$G = \langle x, y \rangle \rtimes A$$

is a polycyclic group. Moreover,  $A$  is self-centralizing and has no cyclic non-trivial  $G$ -invariant subgroups, so that  $G$  is just-infinite. On the other hand, the group  $G$  is complete, i.e. it has trivial centre and  $Aut(G) = Inn(G)$ , and hence  $Aut(G) \simeq G$  (see [9]).

**Lemma 1.** *Let  $G$  be an abelian group. If all proper homomorphic images of the full automorphism group  $Aut(G)$  of  $G$  are finite, then  $Aut(G)$  is finite.*

*Proof.* Assume for a contradiction that  $Aut(G)$  is just-infinite. As the inversion map  $\tau$  of  $G$  belongs to the centre  $Z(Aut(G))$ , we have that  $\langle \tau \rangle$  is a finite normal subgroup of  $Aut(G)$ , so that  $\tau$  is the identity and  $G$  is an infinite abelian group of exponent 2. Let  $\Gamma$  be the set of all automorphisms  $\alpha$  of  $G$  acting trivially on a subgroup of finite index of  $G$ . Then  $\Gamma$  is a non-trivial normal subgroup of  $Aut(G)$  and the index  $|Aut(G) : \Gamma|$  is infinite. This contradiction proves the lemma.  $\square$

**Lemma 2.** *Let  $G$  be a just-infinite group, and let  $N$  be a normal subgroup of  $G$ . Then  $N$  has no finite non-trivial normal subgroups.*

*Proof.* Let  $X$  be any finite normal subgroup of  $N$ . Since  $G/N$  is finite, the conjugacy class of  $X$  in  $G$  is finite, and so it follows from the well known Dietzmann's lemma that the normal closure  $X^G$  is a finite normal subgroup of  $G$ . Therefore  $X = \{1\}$  and the lemma is proved.  $\square$

We can now prove our main result on groups with just-infinite automorphism group.

**Theorem 2.** *Let  $G$  be a group admitting an ascending normal series whose factors are either central or finite. Then the automorphism group  $Aut(G)$  is not just-infinite.*

*Proof.* Assume for a contradiction that the group  $Aut(G)$  is just-infinite, and let

$$\{1\} = G_0 < G_1 < \dots < G_\alpha < \dots < G_\tau = G$$

be an ascending normal series whose infinite factors are central. As the inner automorphism group  $Inn(G)$  is a non-trivial normal subgroup of  $Aut G$ , it follows from Lemma 2 that  $Inn(G)$  has no finite non-trivial normal subgroups. Then the centre  $Z(Inn(G))$  is non-trivial, and so it has finite index in  $Aut(G)$ . In particular, the index  $|G : Z_2(G)|$  is finite, and hence the term  $\gamma_3(G)$  of the

lower central series of  $G$  is finite (see [7] Part 1, p.113). Thus  $\gamma_3(G)$  is contained in  $Z(G)$ , and  $G$  is nilpotent, so that  $\text{Inn}(G)$  lies in the Fitting subgroup of  $\text{Aut}(G)$ . Therefore

$$G/Z(G) \simeq \text{Inn}(G)$$

is a free abelian group of finite rank  $r$  (see [11], Theorem 2). It is well known that the homomorphism group

$$\text{Hom}(G/Z(G), Z(G))$$

is isomorphic to an abelian normal subgroup of  $\text{Aut}(G)$ , so that it is torsion-free and hence also  $Z(G)$  must be torsion-free; moreover,  $\text{Hom}(G/Z(G), Z(G))$  is isomorphic to the direct product of  $r$  copies of  $Z(G)$ . On the other hand, the groups  $\text{Inn}(G)$  and  $\text{Hom}(G/Z(G), Z(G))$  are isomorphic, and hence  $Z(G)$  is infinite cyclic and  $G$  is torsion-free.

Put  $C = Z(G)$  and  $Q = G/C$ , and let  $x$  be any element of  $G \setminus C$ . The mapping

$$\varphi : g \longmapsto [g, x]$$

is a non-trivial homomorphism of  $G$  into  $C$ , whose kernel coincides with the centralizer  $C_G(x)$ , so that  $G/C_G(x)$  is infinite cyclic and

$$G = \langle y \rangle \times C_G(x)$$

for some element  $y$  of infinite order. Let  $m$  be a non-negative integer such that  $(yx)^m$  belongs to  $C_G(x)$ . As  $(yx)^m = y^m x^m z$  for some  $z \in C_G(x)$ , we have that  $y^m$  is in  $C_G(x)$ , and so  $m = 0$ . Therefore  $\langle yx \rangle \cap C_G(x) = \{1\}$ , and hence

$$G = \langle yx \rangle \times C_G(x).$$

It follows that an automorphism  $\alpha$  of  $G$  can be defined by setting

$$y\alpha = yx \quad \text{and} \quad c\alpha = c$$

for all  $c \in C_G(x)$ . Then  $y\alpha^n = yx^n$  for each positive integer  $n$ , so that  $\alpha$  has infinite order and  $\alpha^n$  cannot be an inner automorphism of  $G$ . This is of course a contradiction, since  $\text{Inn}(G)$  has finite index in  $\text{Aut}(G)$ .  $\square$

The above theorem shows in particular that hypercentral groups cannot have just-infinite automorphism groups. We leave here as an open question whether there exists a locally nilpotent group whose automorphism group is just-infinite. As a consequence of Theorem 2, we can observe that the upper central series of any group with just-infinite automorphism group stops at the centre.

**Corollary 1.** *Let  $G$  be a group whose automorphism group  $\text{Aut}(G)$  is just-infinite. Then  $Z(G) = Z_2(G)$ .*

*Proof.* As  $\text{Aut}(G)$  is just-infinite, it follows from Theorem 2 that the index  $|G : Z_2(G)|$  must be infinite, and hence  $Z(G) = Z_2(G)$  because  $Z_2(G)$  is a characteristic subgroup of  $G$ .  $\square$

As infinite simple groups are just-infinite, we have that complete infinite simple groups are trivial examples of groups with just-infinite automorphism groups. Among such groups we find for instance the universal locally finite groups of cardinality  $2^{\aleph_0}$  (see [5]); recall here that a locally finite group  $U$  is said to be *universal* if it contains a copy of any finite group and any two finite isomorphic subgroups of  $U$  are conjugate.

Our last result shows how to find examples of non-simple  $\mathfrak{D}_2$ -groups occurring as full automorphism groups; here the main ingredient is an infinite simple group with finite non-trivial outer automorphism group. Groups of this kind have for instance been constructed by R.J. Thompson in his study of homeomorphisms of the circle  $S^1 = \mathbb{R}/\mathbb{Z}$  (see [2]). Recall here that the *outer automorphism group* of a group  $G$  is the factor group  $\text{Out}(G) = \text{Aut}(G)/\text{Inn}(G)$ .

**Lemma 3.** *Let  $G$  be a group containing a simple normal subgroup  $N$  such that  $G/N$  is finite and  $C_G(N) = \{1\}$ . Then every non-trivial subnormal subgroup of  $G$  has finite index.*

*Proof.* Let  $S$  be any non-trivial subnormal subgroup of  $G$ . Then  $[N, S]$  cannot be trivial, and hence  $S \cap N \neq \{1\}$  (see for instance [10], 13.3.1). Thus  $S$  contains  $N$ , and so the index  $|G : S|$  is finite.  $\square$

**Theorem 3.** *Let  $G$  be an infinite simple group whose outer automorphism group  $\text{Out}(G)$  is finite. Then  $\text{Aut}(G)$  is an infinite complete group whose non-trivial subnormal subgroups have finite index.*

*Proof.* The automorphism group  $\text{Aut}(G)$  is complete by a well known result of Burnside (see for instance [10], 13.5.9), and  $\text{Inn}(G) \simeq G$  is a simple normal subgroup of  $\text{Aut}(G)$  of finite index. Moreover, as  $Z(G) = \{1\}$ , also the centralizer  $C_{\text{Aut}(G)}(\text{Inn}(G))$  is trivial, and hence it follows from Lemma 3 that any non-trivial subnormal subgroup of  $\text{Aut}(G)$  has finite index.  $\square$

## References

- [1] R. BAER: *Finite extensions of abelian groups with minimum condition*, Trans. Amer. Math. Soc. **79** (1955), 521–540.

- [2] M.G. BRIN: *The chamaleon groups of Richard J. Thompson: automorphisms and dynamics*, Inst. Hautes Etudes Sci. Publ. Math. **84** (1996), 5–33.
- [3] F. DE GIOVANNI - A. RUSSO: *The infinite dihedral group as automorphisms group*, Ricerche Mat. **51** (2002), 337–339.
- [4] R.I. GRIGORCHUK: *Just infinite branch groups*, in *New Horizons in pro- $p$  Groups* (edited by M. Du Sautoy, D. Segal and A. Shalev), Birkäuser, Boston (2000), 121–179.
- [5] K. HICKIN: *Complete universal locally finite groups*, Trans. Amer. Math. Soc. **239** (1978), 213–227.
- [6] H.K. IYER: *On solving the equation  $\text{Aut}(X) = G$* , Rocky Mountain J. Math. **9** (1979), 653–670.
- [7] D.J.S. ROBINSON: *Finiteness Conditions and Generalized Soluble Groups*, Springer, Berlin (1972).
- [8] D.J.S. ROBINSON: *Infinite torsion groups as automorphism groups*, Quart. J. Math. (Oxford) **30** (1979), 351–364.
- [9] D.J.S. ROBINSON: *Infinite soluble groups with no outer automorphisms*, Rend. Sem. Mat. Univ. Padova **62**(1980), 281–294.
- [10] D.J.S. ROBINSON: *A Course in the Theory of Groups*, 2nd edition, Springer, Berlin (1996).
- [11] J.S. WILSON: *Groups with every proper quotient finite*, Proc. Cambridge Philos. Soc. **69** (1971), 373–391.