

in characteristic $p > 0$, there is different behaviour; for example, $\mathcal{U}_{2,q}$ has 28 undulations (points where the tangent has 4-point contact). When $g=4$, the curve $\mathcal{C}^6 = \mathcal{F}^3 \cap \mathcal{F}^2$, the intersection of a cubic and a quadric surface, has 60 stalls where the osculating plane meets the curve at four coincident points.

More generally, still with characteristic zero, if \mathcal{C} has genus $g \geq 1$ and $P \in \mathcal{C}$, there exist integers n_1, n_2, \dots, n_g such that no function has pole divisor precisely $n_i P$. Also $\{n_1, n_2, \dots, n_g\} = \{1, 2, \dots, g\}$ for all but a finite number of points. We elaborate this idea and make it more precise in §§8-10.

6. FUNDAMENTAL DEFINITIONS IN ALGEBRAIC GEOMETRY

Let $\mathcal{C} \subset \mathbb{A}^n(K)$ be an irreducible non-singular algebraic curve defined over K , let $I(\mathcal{C}) \subset K[X_1, \dots, X_n]$ be the ideal of polynomials which are zero at all points of \mathcal{C} , let $\Gamma(\mathcal{C}) = K[X_1, \dots, X_n]/I(\mathcal{C})$; and $K(\mathcal{C})$ be the quotient field of $\Gamma(\mathcal{C})$; then $K(\mathcal{C})$ is called the function field of \mathcal{C} . Also, for P in \mathcal{C} let $O_P = \{f/g \mid f, g \in \Gamma, g(P) \neq 0\}$, the local ring of \mathcal{C} at P . Then, by natural inclusions, $K \subset \Gamma(\mathcal{C}) \subset O_P(\mathcal{C}) \subset K(\mathcal{C})$. Also $O_P \setminus \{\text{units}\} = M_P = \langle t \rangle$, the maximal ideal, and for any z in O_P there exist a unique unit u and a unique non-negative integer m such that $z = ut^m$; write $m = \text{ord}_P(z)$. Hence, if $G \in K[X_1, \dots, X_n]$ and g is the image of G in $\Gamma(\mathcal{C})$ with $G(P) \neq 0$, define $\text{ord}_P(G) = \text{ord}_P(g)$. In particular, if \mathcal{C} is a plane curve and $V(L)$ the tangent at P , then $\text{ord}_P(L)$ gives the multiplicity of contact of the tangent with \mathcal{C} .

For the extension of these definitions to the projective case, see Fulton [3], p.182. This is the situation we now consider.

A divisor D on \mathcal{C} is $D = \sum_{P \in \mathcal{C}} n_P P$, $n_P \in \mathbb{Z}$, with $n_P = 0$ for all but a finite number of points P ; the degree of D is $\deg D = \sum n_P$. Then D is effective if $n_P \geq 0$ for all P . For z in $K(\mathcal{C})$, define

$$\begin{aligned} \operatorname{div}(z) &= \sum \operatorname{ord}_P(z) P \\ &= (z)_0 - (z)_\infty, \end{aligned}$$



where

$$(z)_0 = \sum_{\operatorname{ord}(z) > 0} \operatorname{ord}_P(z) P, \text{ the } \underline{\text{divisor of zeros}},$$

and

$$(z)_\infty = \sum_{\operatorname{ord}(z) < 0} - \operatorname{ord}_P(z) P, \text{ the } \underline{\text{divisor of poles}};$$

that is, $\operatorname{div}(z)$ is the difference of two effective divisors and $\deg \operatorname{div}(z) = 0$.

Given $D = \sum n_P P$, define

$$L(D) = \{f \in K(\mathcal{C}) \mid \operatorname{ord}_P(f) \geq -n_P, \forall P\};$$

that is, poles of f are no worse than n_P . In other words, $f \in L(D)$ if $f=0$ or if $\operatorname{div}(f) + D$ is effective.

The set $L(D)$ is a vector space and its dimension is denoted $\ell(D)$.

There is an important equivalence relation on the divisors given by $D \sim D'$ if there exists g in $K(\mathcal{C})$ such that $D - D' = \operatorname{div}(g)$.